

Testing the feasibility & acceptability of selected behaviours to reduce child exposure to indoor air pollution in rural South Africa

Brendon Barnes
Angela Mathee

May 2003

Health and Development Research Group
Medical Research Council of South Africa



**BUILDING A
HEALTHY NATION
THROUGH RESEARCH**

**ENVIRONMENT AND
DEVELOPMENT**

**NATIONAL
PROGRAMME**

[INSIDE FRONT COVER]

Published by:

The Medical Research Council of South Africa,
Health & Development Research Group,
Environmental Health Office
Johannesburg
<http://www.mrc.ac.za/healthdevelop/healthdevelop.htm>

ISBN: 1-919809-65-1

Suggested citation:

Barnes, BR & Mathee A. Testing the feasibility and acceptability of selected behaviours to reduce child exposure to indoor air pollution in rural South Africa. Johannesburg: Medical Research Council of South Africa. 2003.

Copies:

Copies of the report can be obtained from:
The Medical Research Council of South Africa,
Health & Development Research Group,
Environmental Health Office
P. O. Box 87373, Houghton, 2041
Johannesburg, South Africa

Phone: (011) 643 7403
Fax: (011) 642 6832
Email: bbarnes@mrc.ac.za

TABLE OF CONTENTS

ACKNOWLEDGEMENTS	V
ACRONYMS	VI
EXECUTIVE SUMMARY	1
1. BACKGROUND.....	4
2. STUDY SETTING.....	6
3. PHASE TWO METHODOLOGY	7
3.1 Study design	7
3.2 Research participants	7
3.3 Procedure.....	8
3.4 Personnel	10
3.5 Analysis.....	10
4. RESULTS	10
4.1 Impact on behaviours	10
4.2 Perceived benefits and sustainability of behaviour change	20
5. DISCUSSION & RECOMMENDATIONS	21

FIGURES, TABLES & BOXES

Figure 1 Phases of the project.....	5
Figure 2 Mixing a paste and sealing a leaking stove	11
Figure 3 Methods of sealing chimneys	11
Figure 4 Improvements in ventilation practices (n=20).....	13
Figure 5 Non-functioning windows that have been covered.....	13
Figure 6 Improvements in child location practices (n=16).....	14
Figure 7 Number of households that improved child location practices by presence of an extra caregiver	15
Figure 8 Reductions in the duration of solid fuel burning (n=13)	17
Figure 9 Perceived benefits of behaviour change	20
Table 1 Breakdown of research participants	7
Table 2 Dates of phase two activities	9
Table 3 Summary of behaviours	19
Box 1 Negotiating behaviours with families	9

ACKNOWLEDGEMENTS

Funding for this project from the United States Agency for International Development through the CHANGE project (implemented by the Academy for Educational Development with the Manoff Group) and the South African Medical Research Council is gratefully acknowledged. The authors would also like to Dr. Laurie Krieger, Lonna Shafritz, Mike Favin and Lisa Sherburne (staff members of the CHANGE project, the Academy for Educational Development and the Manoff Group) for their valuable technical contribution to the project. The assistance of the following researchers - Kebitsamang Moilola, Charity Masilo, Brenda Mogokonyane and Valencia Bonnyane - is also gratefully acknowledged.



ACRONYMS

ALRI	Acute lower respiratory infections
CO	Carbon monoxide
COLD	Chronic obstructive lung disease
EHP	Environmental Health Project
LPG	Liquid petroleum gas
MRC	Medical Research Council of South Africa
NO ₂	Nitrogen dioxide
PM _{2.5}	Particulate matter of 2.5 µm or less
RSP	Respirable suspended particulates
SO ₂	Sulphur dioxide
SPSS	Statistical Package for the Social Sciences
TIPs	Trials of Improved Practices
USAID	United States Agency for International Development

EXECUTIVE SUMMARY

Background: Indoor air pollution has been causally linked to Acute Lower Respiratory Infections in children less than five years old and accounts for a significant proportion of death and illness in developing countries. At the level of prevention, behaviour change has been identified as a possible intervention strategy to reduce the impact of indoor air pollution on child respiratory health. Yet published studies have yet to systematically focus on the effectiveness of behaviour change strategies in reducing child exposure to indoor air pollution.

The aim of this study, therefore, was to test the acceptability and feasibility of four clusters of behaviours: 1) improve stove maintenance practices, 2) improve ventilation practices, 3) improve child location practices & 4) reduce the duration of solid fuel burning. These behaviours were identified based on a previous phase of research, which aimed to identify indoor air pollution related behaviours that were protective of child respiratory health.

Methods: The study utilised a Trials of Improved Practices methodology, which is a formative research method that involves families actually trying out and possibly modifying selected behaviours over a selected period of time. Thirty families were selected to participate, all of who agreed to try one or more of the above-mentioned behaviours over a four-week trial period. Using participatory observation techniques, the behaviours were assessed before and after home visits by researchers during winter (July-August) 2002. The final visit also involved in-depth interviews to gain an understanding of the barriers, motivations and sustainability of the behaviours.

Results: Of the 30 families that participated in the study, fixing of stoves was recommended to 15. Eleven of the 15 families agreed to attempt the behaviour(s). However, fixing wood stoves proved to be difficult for the majority of these households. Only three out of the 11 families managed to fix their stoves over the trial period. Motivations for fixing stoves included improvements in the levels of air quality, reductions in the amount of dust and soot in the house and generally a cleaner house. In addition, mothers reported that their houses smelled better. The cost of buying materials and hiring someone to fix them was cited as the main barrier to fixing up stoves.

Opening at least two sources of ventilation for longer periods of time while a fire was burning was recommended to 29 families. All 29 families agreed to attempt the behaviour(s) over the trial period. Twenty of the 29 families managed to improve their ventilation practices. The percentage of time that two sources of ventilation were opened while a fire was burning was used as an indicator for ventilation practices. For morning burnings, ventilation practices improved from 7% to 19% amongst families that improved their practices. For evening burnings, ventilation practices

improved from 10% to 23%. Motivations for opening ventilation included improvements in the levels of air quality, reductions in the amount of dust and soot in the house and a cleaner smelling house. Non-functioning windows and the cold weather were identified as barriers to improving ventilation practices.

Reducing the amount of time that their children spend close to a burning fire was recommended to 28 families. All 28 families agreed to try this. At the end of the trial, 16 of the 28 households successfully managed to do this. The percentage of time that children were in close proximity to the stove while a fire was burning was used as an indicator for child location practices. The average proportion of time that children were close to the stoves while a fire was burning was reduced from 43% to 29% amongst the families that improved this behaviour. During evening burnings, the average figure was reduced from 41% to 27%. The 16 mothers who successfully improved their child location practices cited the fact that their children's lungs were protected from smoke as the main motivation for moving them away from the fire. Moreover, four of the 16 mothers reported that it was easier to do chores in the kitchen while someone else was looking after their children. Barriers to keeping children away from fires include the fact that many children did not like being away from their mothers and that mothers felt their children would be cold by being away from the fire. Mothers who did not have someone else to assist them with looking after their children allowed their children to play in another room with other children.

Reducing the length of solid fuel burning was recommended to 27 families. Twenty-six families agreed to attempt this. Only 13 managed to reduce the length of solid fuel burning over the trial period. During morning burnings, the average length of solid fuel burning was reduced from 250 minutes to 219 minutes amongst families who reduced the duration of solid fuel burning. Slightly greater reductions were achieved for evening burnings, where the duration of burning was reduced from 242 to 198 minutes. Of the 13 families that managed to reduce the duration of solid fuel burning, five indicated that they saved fuel by doing so. Burning for shorter periods of time resulted in monetary savings from buying less wood. In addition, this reduced the time & effort required to collect cow dung (which is freely available). Eight families indicated that their motivation for reducing the duration of solid fuel burning was that their children were less exposed to smoke over a prolonged period of time. The need for a fire to be burning for prolonged periods to be able to perform domestic chores, as well as the cold weather, was cited as the main barriers to reducing the duration of solid fuel burning.

Half of all mothers (n=15) reported improvements in their children's respiratory health as a result of behaviour change. Eleven families (36%) reported that they benefited by having a cleaner and

less smelly house. Fixing up their stoves, opening windows and doors as well as reducing the amount of time that stoves were burning resulted in a reduction of the amount of dust and soot accumulating on hard surfaces and curtains in the kitchen. This, it was reported, made cleaning the house, particularly the kitchen a little easier to do. The house also did not smell as much of smoke. Five families (17%) reported that they saved fuels (and consequently money to buy fuels) by reducing the duration that solid fires were left to burn. Of the 30 households that participated in the study, 25 (83%) indicated that they will continue and 5 (17%) reported that they would not continue with the behaviours as they were too difficult to perform.

1. BACKGROUND

Estimates indicate that indoor air pollution is responsible for between 2.7 and 2.8 million deaths annually (Bruce, Perez-Padilla & Albalak, 2000). In many developing countries, indoor air pollution accounts for as much as 4-6 percent of the burden of disease, placing it above environmental tobacco smoking, sexually transmitted diseases, alcohol and homicides as a leading cause of ill health and death (Smith, 1999). Two recent reviews of published epidemiological studies have identified Acute Lower Respiratory Infections (ALRI), such as pneumonia, amongst children under 5 years of age in developing countries, as one of the key health outcomes of exposure to indoor air pollutionⁱ (Smith, Samet, Romieu & Bruce, 2000, Bruce *et al.*, 2000).

Indoor air pollution in developing countries mostly arises from the indoor burning of solid fuels such as wood, animal dung, coal and crop residues in open fires or poorly functioning stoves. The incomplete combustion of these fuels releases pollutants such as particulate matter (PM), carbon monoxide (CO), sulphur dioxide (SO₂), nitrogen dioxide (NO₂) and other organic compounds into the living environment (Smith, 1987). It is estimated that at least two-thirds of all households in developing countries are still primarily dependent on biomass fuels and coal, affecting approximately 2 to 3.5 billion people worldwide (The World Resources Institute, 1998).


Other determinants of poor indoor air quality include the stoves that people use (or lack thereof) (Ezzati & Kammen, 2002a, Albalak *et al.*, 2001 & Wafula *et al.*, 2000) and ventilation (Nystrom, 1994 in Ballard-Tremmer & Mathee, 2000, Gitonga, 2001). While air quality is important, the health effects of indoor air pollution are also determined by the amount of time a person spends breathing the polluted air. Studies have shown that people in developing countries spend between three and seven hours a day in the burning environment (Bruce *et al.*, 2000). Because of their domestic responsibilities in and around the burning room, women are typically exposed to indoor air pollution far more than their male counterparts (Engel, Hurtado & Ruel, 1998). In addition, women usually have the added responsibility of caring for children. As a result children tend to be close to their mothers - either carried on their mothers' backs or within eyesight – thus also exposing them to high levels of indoor air pollution on a daily basis.

At the level of prevention, behaviour change has been identified (amongst others) as a possible intervention strategy to reduce the impact of indoor air pollution on child health. Behaviours such as moving children out of the room while a fire is burning, using pot lids while cooking and improving the quality of ventilation practices have all been identified for their potential to reduce the impact of indoor air pollution on human health (cf. von Schirnding, Bruce, Ballard-Tremmer,

Ezzati & Lvovsky, 2002). Yet published research studies have yet to systematically focus on these and other behavioural determinants of exposure to indoor air pollution or the effect of behaviour change strategies in reducing the impact of indoor air pollution (Ezzati & Kammen, 2002a, Barnes & Mathee, 2002a & Favin, Yacoob & Bendahmane, 1999).

In response to this, a programme of work is underway in rural South Africa to evaluate the effectiveness of such an intervention. Overall, the project is divided into four phases.

Figure 1 Phases of the project



Phase one identified and recommended four behavioural clusters that may serve to reduce child exposure to indoor air pollution: 1) stove maintenance, 2) increase the duration that two sources of ventilation are opened during burning, 3) reduce the duration of burning and 4) reduce the amount of time that children spend in close proximity to the stove. To identify these behaviours, the behavioural patterns of 20 families that cared for a child with a history of severe ALRI were observed and compared with the behavioural patterns of 20 families that cared for a child with normal respiratory function. In addition, mothers were interviewed after the observations and a further 27 mothers participated in focus group discussions to identify recommendations to reduce children's exposure to indoor air pollution. The four behavioural clusters were selected based on their association with the risk of respiratory ill health, suggestions from mothers, likely cost, required effort and probable reductions in exposure to indoor air pollution (cf. Barnes & Mathee, 2002b).

This report highlights the findings of phase two, the goal of which was to test the feasibility and acceptability of the four behavioural clusters mentioned above. To do this, research attempted to answer the following questions:

- To what extent were mothers willing and able to perform the behaviours?
- How did they perform the behaviours?
- What were the barriers to the performance of the behaviours?
- What were the motivating factors for implementing the behaviours?
- Did mothers perceive any impact of behaviour change?
- Will families continue to perform the behaviours after the trial period?

2. STUDY SETTING

Research participants were residents of Goedgevonden and Uitkyk – two poor, rural villages in the North West Province of South Africa. The villages are situated within 20 kilometres of the villages used in phase one. Dwellings consist of two to four rooms, one of which normally serves as the kitchen (which is also the room in which burning takes place). Ventilation, in the form of windows and a door leading to the outside, are available in all households. However, in many instances, these are in poor condition and are not functioning properly (see barriers to ventilation practices below).

People living in these communities are extremely poor and have high unemployment rates. Households are dependant on wood, cow dung, crop residues and kerosene for their energy requirements. During winter, solid fuels are typically burned in poorly maintained wood stoves, which are normally situated in the kitchen. Kerosene is burned in kerosene stoves. For example, indoor air quality measurements taken during July 2001 (winter) showed very high concentrations of particulate matter (PM_{2.5} and RSP) (Barnes & Mathee, 2002b).

The area becomes extremely cold during winter (average winter minimum temperature is 5°C but temperatures often fall below 0°C) and because of the need for space heating, this is the time of the year when most indoor burning takes place. Fires are typically ignited early in the morning and left to burn out after cooking and water heating is completed. Fires are re-ignited in the late afternoon to cook the evening meal and left to burn out for space heating. Summer cooking practices typically involve cooking outside on an open fire (or a brazier) or using kerosene stoves (Mathee *et al.*, 2000).

Ventilation practices typically involve opening a source of ventilation such as a window or a door for 15 minutes during ignition or until visible smoke disappears. Ventilation is then closed for the

rest of burning. The opening of at least two sources of ventilation was found to be significantly associated with improved child respiratory health in phase one. Importantly, young children often follow their mothers around and can spend up to 6 hours close to a burning fire during winter. Older children are normally left to sit close to the fire while younger children are carried on their mothers' backs. Having the child away from the burning fire was found to be associated with lower risk of child respiratory illness in phase one (Barnes & Mathee, 2002b).

3. PHASE TWO METHODOLOGY

3.1 Study design

Phase two utilised a Trials of Improved Practices (TIPs) (Dickin & Griffiths, 1997), which is a formative research method that involves families actually trying out and possibly modifying selected behaviours over a selected period of time. Thirty households were selected to participate in the study. Behaviours were assessed before and after home visits by researchers during winter (July – August) 2002. This technique allowed an analysis of the acceptability (willingness to try) as well as the feasibility (ability to perform) of the behaviours.

3.2 Research participants

Research participants were 30 mother-child combinations divided into groups based on pre-selected criteria that were found in phase one to have an impact on the behaviours being tested - for example, the availability of an adult to assist the mother by looking after the child in a location away from the burning fire, the age of the study child and the type of appliance used (Barnes & Mathee, 2002b).

Table 1 Breakdown of research participants

	Have used wood stove in the last two weeks				Have not used wood stove in the last two weeks.
	Children 1-5 years		Children <1 year old		
	Adult present to watch child in home during burning	No adult to watch child in home during burning	Adult present to watch child in home during burning	No adult to watch child in home other than mother	
Goedgevonden:	3	3	4	4	1
Uitkyk:	3	3	4	4	1
Sub-Totals	6	6	8	8	2
Total	30				

3.3 Procedure

Research participants were identified based on door-to-door visits to approximately 150 households by researchers. During these visits, researchers administered a screening questionnaire that included questions on the availability of an additional adult to look after the children while a fire was burning, the age(s) of children in the home and whether or not a wood stove was used in the previous two weeks. Once selected, households were informed about the study and then asked to participate and to give informed consent. All households who were selected, agreed to participate in the study. Dates and times were agreed upon for researchers to assess behaviours.

Behaviours were assessed during the week of 3-11 July 2002 when minimum temperatures first dropped below 5°C (assessment visit). Typically, researchers would arrive at a household at approximately 06h30 (or in time for the first burning) and observe household behaviours for a 12-hour period while filling in a semi-structured observation sheet. Where necessary, they would ask and respond to mothers' questions and participate in household activities. After each day, a team meeting was held to formulate recommendations for each household. Recommendations were based on observed behaviours and tailored to each family's need(s). In some cases only one behaviour was recommended while in others all four behaviours were recommended.

Researchers visited each home to offer the recommendations to mothers and family members. All available family members were invited to join the discussion. During this counselling visit researchers firstly explained (in the seTswana language) how the fuels that people burn produces smoke that contains dangerous chemicals and that breathing in this smoke affects their family's health. Although researchers offered selected behaviours, no recommendations were forced upon families. Instead, researchers assisted each family, through a process of negotiation, with finding practices that it felt would be most feasible. In many instances families felt that, from the outset, certain behaviours would be too difficult to perform.

Once family members decided which behaviours they would try, researchers then facilitated a discussion of *how* they would perform those behaviours. Families were asked questions such as: who is going to take responsibility for looking after the child while the mother was cooking? Do you have enough clothes to keep the child warm while away from the fire? If the mother can look after the child away from the fire, can someone else do her chores? Who is going to fix the stove? How are you going to fix it? How much money do you need for this? If you reduce the length of time that a fire is burning, how will you do all of your household chores in such a short period of time? I see that your windows cannot open, are you able fix them? Who will do this? How much money do you need for this?

Box 1 Negotiating behaviours with families

Mpho is the mother of a 5-month old daughter and lives with her sister Katlego. Typically Mpho starts a fire using wood and cow dung in her wood stove at 06h00, opens one window for 10 minutes during ignition and then closes it for the rest of burning which ends at about 09h00. When her daughter wakes up at about 07h00, she is left in a cot next to the stove while Mpho performs her domestic chores.

During the counselling visit, the researcher discussed the dangers of indoor air pollution with Mpho and Katlego and recommended that they attempt to open the main kitchen window for 35 minutes and also open an additional window/door for the same length of time. Mpho believed that they would not be able to open two windows for 35 minutes because of the cold. They negotiated a time that would be suitable for them. At the end of this process, it was agreed that they would open at least one window for 25 minutes and the other for 10 minutes (or at least until the smoke from ignition process had cleared).

The researcher also recommended that the child be moved away from the stove to a room that has a door that could be closed for the entire duration of burning. Katlego agreed to look after the child in that room while a fire was burning but indicated that she could not do this for the entire time that the fire was burning because she also has to care for herself e.g. bathe and perform domestic chores like washing dishes. A suitable time was negotiated with Katlego when she could bring the child into the kitchen – particularly during periods of low emissions. She also agreed that, when in the kitchen, the child should be left in the cot as far away from the stove as possible. In this way, behaviours were negotiated with family members and solutions were tailored, not only to their needs, but also to what they were realistically able to achieve.

After two weeks researchers visited each family to encourage them to continue with the behaviours for the rest of the trial period (reminder visit). After another two weeks, researchers visited each household to assess the behaviours as well as conduct a semi-structured interview with mothers. During this interview, mothers were asked questions such as: Were you able to perform the behaviour(s)? If no, what were the reasons? If yes, how easy was it? What made it enjoyable? Was there anything you did not like? Will you continue to do this? Will you need to change anything to be able to continue? If you were to stop, what would make you stop? In this way, the study could identify the feasibility of each behaviour specific to each household, the barriers as well as the motivations of the behaviours.

Table 2 Dates of phase two activities

Activity	Dates (2002)
Assessment visits	07/03– 07/11
Counselling visits	07/14 – 07/18
Reminder visits	08/04 – 08/06
Final visit	08/18 – 08/23

3.4 Personnel

The project involved four researchers (Kebitsamang Moilola, Charity Masilo, Brenda Mogokonyane and Valencia Bonnyane), a project co-ordinator (Brendon Barnes) and a project manager (Angie Mathee). Researchers participated in a five-day training course (07/24-28), co-ordinated by Brendon Barnes, in Mafikeng before the study commenced. The course involved training in: 1) principles of TIPs, 2) indoor air pollution and health, 3) qualitative interviewing, 4) observation techniques, 5) note taking and 6) counselling skills. In addition, the research instruments and researchers' skills were tested and refined during a two-day pilot phase in Bethel, a village situated 20 kilometres south of the study villages.

3.5 Analysis

Data from the observations of household behavioural patterns were captured and analysed using the Statistics Package for the Social Sciences (SPSS) software package. To observe overall trends in the data, frequency distributions and measures of central tendency were used. Qualitative data were analysed using a thematic analysis (Miles & Huberman, 1994).

4. RESULTS

4.1 Impact on behaviours

Stove maintenance practices

Depending on the stove problem, researchers negotiated with families to either seal their stoves and chimneys with a product such as putty (readily available at the local store) or replace missing parts. If cheaper alternatives that would suit families' circumstances and budgets were available, researchers assisted families to identify them.

Of the 30 families who participated in the study, it was recommended that 15 fix their wood stoves. Eleven families agreed to do this. The four families that did not agree to fix their stoves cited the cost of fixing stoves as the primary reason for not attempting the behaviour(s). Fixing wood stoves proved to be very difficult for the majority of households: only 3 out of the 11 families managed to fix their stoves.

The three households that fixed their stoves and chimneys used a variety of low cost methods to do so. These included mixing a paste of cow dung, soil, wool (as a binding agent) and water to seal leaking parts of stoves (see figure 2). One household used a mixture of cement, coarse salt and water as a sealant. To fix chimneys, families either tied an old cloth or fixed an old tin sheet over the broken part of the chimney (see figure 3). Although these methods are relatively

temporary, they are however relatively easy to do, cheap and can be done quickly. More importantly, observations and reports from mothers show that they are effective in reducing the emissions from stoves. According to one mother, *“At first I was not going to fix my stove because I did not have enough money for it. I did not know about mixing cow dung and water until a friend’s mother told me about it. It seems to work so I’ll keep on doing it.”*

Figure 2 Mixing a paste and sealing a leaking stove



Figure 3 Methods of sealing chimneys



Motivations for fixing stoves reported by the three families that fixed their stoves included improvements in the levels of air quality, reductions in the amount of dust and soot in the house and generally a cleaner house. This made cleaning of hard surfaces and curtains easier to do. In addition, after fixing their stoves, mothers reported that their homes smelled better.

For the eight families that did not fix their stoves, the monetary cost, not only of purchasing materials such as putty or tin, but also of hiring someone to do it was thought to be prohibitive.

Many families did not have a person who had the skills to fix stoves and therefore had to pay a member of the community to do it. Households that did not have someone living in the house to fix up the stove and did not have the money to pay someone, usually had to rely on the goodwill of friends and neighbours. Very often, this meant that they would have to wait until that person was available. For example, in one household, the person who was meant to fix their stove was ill, while in another the person was too busy and as such, no improvements were made to the stove over the trial period.


It should be noted however, that even though the compliance rate was low for fixing stoves, mothers who could not fix their stoves nevertheless attempted various behaviours on their own that revolved around the stove but did not necessarily relate to what they agreed to do. These included cleaning the chimney, replacing missing stove lids with a pot and drying wood and cow dung before burning it. As one mother put it, *“Even though I didn’t have money to fix the stove, your visit reminded me to at least clean my chimney with an orange sack and a stick. This was easier to do because I do this all the time.”*

Ventilation practices

Household members, with the assistance of researchers, agreed to and set themselves goals for opening two sources of ventilation during burning. Overall, improving the quality of double ventilation practices was recommended to 29 families, all of whom agreed to do so. Four of the 29 families also agreed to fix broken windows in the room used for burning. Twenty of the 29 families managed to improve their ventilation practices over the trial period.

The percentage of time that two sources of ventilation were opened while a fire was burning was used as an indicator of ventilation practices. Amongst the 20 families that improved this practice, the proportion of time that two sources of ventilation were opened while a fire was burning increased from 7% to 19% during morning burnings. For evening burnings, this figure improved from 10% to 23%.

Figure 4 Improvements in ventilation practices (n=20)



The eleven families that improved their double ventilation behaviours cited a number of motivations for doing so. These included the presence of less smoke in the house while a fire was burning, hard surfaces were easier to clean because less soot was deposited in the living environment and their homes generally smelled better. A number of barriers to improving ventilation practices were identified, most notably, non-functioning windows in the room used for burning. Usually both the windowpane and the opening mechanisms were broken. To keep out the cold as well as to secure privacy, people normally covered windows with water-resistant plastic bags, cardboard or a sheet of metal. Figure 5 shows two of the study dwellings with non-functioning windows that have been covered with plastic and metal sheeting. The cost of fixing windows was thought to be too high, and as a result, none of the four families that agreed to fix broken windows did so.

Figure 5 Non-functioning windows that have been covered




Another barrier to opening ventilation was the cold winter temperatures. At least 4 out of the 9 households who agreed to but did not improve their ventilation practices cited cold weather as the primary reason for doing so. In these cases the door leading to the outside was usually opened only for a short period during ignition and then closed again for the rest of the burning. As one mother put it, *“No, I did not open my windows because it was too cold, I only opened the outside door for a short time thinking that it is enough for the smoke to go through.”*

Reducing the amount of time that children spend close to the fire

Mothers were asked to reduce the amount of time that their children spend in close proximity to the stove during burning. Overall, it was recommended that 28 families reduce the length(s) of time that their children were close to fires. All 28 families agreed to try this for the trial period. Sixteen of the 28 families successfully managed to reduce the length (s) of time that their children were in close proximity to the stove. The percentage of time that children were in close proximity (within 1.5 metres) to the stove while a fire was burning was used as an indicator for child location practices. The proportion of time that children were close to the stove while a fire was burning was reduced from 43% to 29% during morning burnings amongst the 16 families that improved this behaviour. During evening burnings, the figure was reduced from 41% to 27%.

Figure 6 Improvements in child location practices (n=16)




Even though keeping children away from the burning fire was reportedly difficult, the 16 mothers who were able to do so cited the fact that their children were not exposed to indoor air pollution as the main motivation for doing so. The reassurance that they were protecting their children’s respiratory health was a significant motivating factor for persisting with the behaviour. Moreover,

four of the 16 mothers reported that it was easier to do chores in the kitchen while someone else was looking after their children.

Mothers reported a number of barriers to moving their children out of the burning room. Many children were reportedly not used to being looked after by someone else and were often unhappy to be away from their mother and resorted to crying and misbehaving. After a while, mothers allowed their children to be close to them and to the burning fire. This is highlighted in the following extract taken from an interview with a mother, “ *I felt very bad because my child would cry. I did not want to see him cry because he would only want to be with his mummy and not with other people in the house. He only wanted to be with me.*” It was also cold being away from the fire, so many mothers felt that it was cruel and un-nurturing to allow their children to be cold while a fire was burning.

As an important behaviour in terms of exposure to indoor air pollution, and in light of the finding in phase one that the availability of someone to look after the child in a location away from the stove was found to be a significant determinant of child location practices, it was decided to classify families based on the presence or absence of an additional adult to look after the child. No differences were evident between households with and without extra adults and whether or not they managed to improve child location practices. Both groups managed to reduce the amount of time that their children spent close to the burning fire by 14% during morning burnings and by 13% during evening burnings. Interestingly, those households without extra caregivers were slightly over represented in the group that reduced the amount of time that their children spent in close proximity to the burning fire.

Figure 7 Number of households that improved child location practices by presence of an extra caregiver




How did mothers without the assistance of other adults manage to keep their children out of the room while a fire was burning? Of the eight mothers who managed to do this, six left their children to play with other siblings (usually younger than five years old) in another room of the house while a fire was burning. Typically, mothers in this group would give them toys to play with and dress them warmly so that they would not feel cold. They would also instruct the oldest sibling to take care of the younger children. These children were typically left unsupervised for this duration.

One mother (of a two year old son) without additional assistance, and without any other children, attempted herself to spend less time in proximity to the fire. *"In order to keep him away, I avoided staying in the burning room because every time I am in there, he would be holding my dress. When we were talking before it sounded simple, but it was difficult because this delayed my daily duties because the child follows me wherever I go. I could cook freely only when he was sleeping so I had to wait for him to sleep. At the same time my pots would be burning because there is nobody to look after him."* While a fire was burning, one mother (without additional assistance) took her child to her mother who lived two dwellings from her.

Duration of burning

Results from phase one showed that many households did not extinguish fires once cooking and water heating was completed and the house was sufficiently warm. Instead, fires were left to extinguish on their own, which could take up to two hours after cooking and water heating were completed. Families were asked to reduce the amount of time that solid fuels were burned, particularly after cooking and water and space heating were completed.

Overall, it was recommended that 27 families reduce the duration of solid fuel burning of which 26 families agreed to try for the trial period. Only 13 of the 26 families managed to successfully reduce the length of solid fuel burning over the trial period. During morning burnings, the average length of solid fuel burning was reduced from 250 minutes to 219 minutes amongst the 13 families that improved this behaviour. Slightly greater reductions were achieved for evening burnings: the duration of solid burning was reduced from 242 to 198 minutes. Extinguishing a fire normally entailed pouring water or leftover maize porridge over the embers to extinguish them or using just enough fuel as is necessary to complete the cooking, water and space heating.

Figure 8 Reductions in the duration of solid fuel burning (n=13)

Of the 13 families that managed to reduce the duration of solid fuel burning, five indicated that they saved fuel by doing so. Burning for shorter periods of time resulted in monetary savings from buying wood. In addition, using less fuel reduced the time & effort required to collect cow dung, which is freely available. Eight families indicated that their motivation for reducing the duration of solid fuel burning was that their children were less exposed to smoke over a prolonged period of time.

A number of barriers to reducing the duration of solid fuel burning were evident, most notably the need for warmth during the cold winter months. Many mothers reported that when they extinguished their fires, their homes would become extremely cold – particularly during mornings and evenings. In addition, reducing the length of time that a fire was burning often meant that the mother had to do all the chores that rely on the fire during a shorter period of time. These chores include heating water for tea, heating babies' bottles, bathing and reheating leftover food. This was often very difficult to do because the fire would be needed for different purposes at different times.

Box 2 The burden of reducing the duration of burning

Wilhelminah, a mother of three, lives in a three-roomed dwelling. She lives with her mother-in-law and children except when her husband, who works in Mafikeng, comes home every second weekend. Her three children are aged 9 years, 7 years and 7 months. The two eldest children are at school, while the youngest stays at home with her during the day. Wilhelminah typically gets up at about 05h45 and starts a wood fire at 06h00. She uses the fire to heat leftover food for breakfast and begins to heat water on her wood stove.

She wakes her two older children up at 06h10. After getting out of bed, they stand next to the fire to heat themselves, use the warm water to bathe with, eat and then leave for school at about 07h00. At 08h00, she heats more water to wash dishes with and then starts cooking. Her youngest child wakes up a little after 08h00, after which she heats more water to warm her baby's feeding bottle. She changes her baby's napkin and then heats more water to bathe her child with at about 08h35. She also uses this hot water to wash her child's napkins with at about 09h00. She continues to cook until 10h00. She uses the fire to heat her iron with and then irons her family's clothing with until about 10h30. After four-and-a-half hours, the fire is left to extinguish on its own. By this time, she has also already fetched wood from outside a few times, collected dried cow dung (as her wood supplies are running low), collected fresh water from the village well, and has begun to wash her family's clothing. While being used continuously from 06h30 until 10h30 for domestic tasks, the fire simultaneously provides comforting heat during cold winter mornings.

In short, having a fire burning for prolonged periods of time allows mothers to do the chores that rely on heat when they need to be done while simultaneously providing much needed warmth for the household. Reducing the length of burning would entail having to do all of those chores during a shorter time period, or alternatively igniting a fire each time a chore needs to be done.

Table 3 summarizes the impact of the home visits on the four clusters of behaviours.

Table 3 Summary of behaviours


Behavioural cluster	Number of families to which recommended.	Number of families that declined to perform the behaviours.	Number of families that agreed to perform the behaviours.	Number of families that successfully performed the behaviours.	Description of how behaviours were performed.	Barriers	Motivations
Improve stove maintenance practices.	15	4	11	3	Families used a mixture of cow dung, soil and water to seal leaking parts of stoves and used old cloths or old tin sheets to cover broken parts of chimneys.	<ul style="list-style-type: none"> ▪ Cost of purchasing materials. ▪ Cost of hiring person to fix stoves. 	<ul style="list-style-type: none"> ▪ Less smoke in he house. ▪ Less dust which makes cleaning easier. ▪ Cleaner smelling house.
Improve double ventilation practices.	29	0	29	20	Families opened doors and windows for longer periods of time while a fire was burning.	<ul style="list-style-type: none"> ▪ Non-functioning windows that cannot be opened. ▪ Cold weather. 	<ul style="list-style-type: none"> ▪ Less smoke in the house. ▪ Less dust which makes cleaning easier. ▪ Cleaner smelling house.
Improve child location practices in relation to the fire.	28	0	28	16	Families attempted to keep children out of the kitchen while a fire was burning. To do this, children were looked after by another adult or left to play with other children in another room of the house.	<ul style="list-style-type: none"> ▪ Children often do not like being away from their mothers. ▪ Lack of extra adult supervision for children when they are away from the kitchen. ▪ Children become cold when out of the kitchen 	<ul style="list-style-type: none"> ▪ Knowledge that children's respiratory health is protected.
Reduce the duration of solid fuel burning	27	1	26	13	Families reduced the length of solid fuel burning by extinguishing the fire after cooking and water and space heating was complete.	<ul style="list-style-type: none"> ▪ House becomes cold when fire is extinguished. ▪ Increased burden on women to perform chores in shorter time. 	<ul style="list-style-type: none"> ▪ Fuel saving. ▪ Less money needed to purchase wood. ▪ Less time & effort required to collect cow dung.

4.2 Perceived benefits and sustainability of behaviour change

In line with the motivations cited above, mothers noted what they perceived to be benefits of behaviour change over the trial period. Half of all mothers (n=15) reported improvements in their children’s respiratory health following the trials. The reported improvements ranged from a decrease in specific symptoms, such as coughs, to less frequent visits to the local hospital.

Eleven mothers reported that they benefited by having a cleaner and less smelly house. Fixing up their stoves, opening windows and doors, as well reducing the amount of time that stoves were burning resulted in reductions in the amount of dust and soot that accumulated on hard surfaces and curtains in the kitchen. This, it was reported, made cleaning the house, particularly the kitchen, a little easier to do. The house also did not smell as much of smoke. Only five families reported that they saved fuels and consequently money to buy fuels as well as effort in collecting free fuels such as cow dung by reducing the time that solid fires were left to burn.

Figure 9 Perceived benefits of behaviour change



Of the 30 families that participated in the study, 25 (83%) reported that they intended to continue with the behaviours, mainly due to improvements in their children’s health. Five (17%) families reported that they would not continue with the behaviours because they were too difficult to perform.

5. DISCUSSION & RECOMMENDATIONS

Behaviour change is notoriously difficult to achieve and even more difficult to sustain. A recent review of published behaviour change studies found that only a small percentage of studies managed to achieve behaviour change and those that did, were plagued with methodological flaws (Cave & Curtis, 1999). This report showed how, after home visits to discuss indoor air pollution and behaviour change, families changed their behaviours (to varying degrees) to reduce their children's exposure to indoor air pollution. However, it is important to highlight the methodological weaknesses of the approach used in this study.

Firstly, it is possible that people's behaviours were influenced by the presence of a researcher. During the follow-up visits families were already sensitised to the fact that researchers were there to determine whether they had performed the agreed behaviours. Consequently, the behaviour change reported here could be a result of *reactivity* due to the presence of researchers and not true behaviour change. The same weakness holds true for the personal interview conducted during the follow-up visit. The study attempted to circumvent this by stressing to families that the researchers would learn just as much from what they were not able to do as from what they were able to do. They were therefore encouraged to be as open and honest about what they were and were not able to perform.

Secondly, because there were only two data collection visits (one before and one after), it was not possible to capture important daily and weekly variability in behaviour. These two visits represent mere 'snapshots' in the lives of the families that participated in the study. For example, it is not known how a particularly cold spell would affect families' abilities to open windows, keep their children away from the stove and reduce the time that a fire was burning. Moreover, factors such as how cold it was on the assessment days and whether someone else was present on that particular day to look after the child would all influence the behaviours that were observed.

Nevertheless, the study yielded valuable information with regard to behaviour change and indoor air pollution. It showed how indoor air pollution behaviours are firmly rooted in poverty, weather and an enabling environment. Importantly, it showed how certain behaviours such as ventilation and child location practices were easier to improve than stove maintenance and reducing the length of burning.

For example, the cost of fixing of wood stoves was found to be prohibitive for most families that participated in the trials. The three families that did fix their stoves and chimneys used relatively

low cost methods to do so. The fact that so few families managed to fix their stoves indicates that the promotion of stove fixing may only be feasible for a small number of people in this context. In addition, even though families reported improved levels of air quality, observations indicate that 'fixed' stoves were still emitting visible smoke albeit to a lesser extent than before. Because of the perception that stoves were not emitting pollutants, it was found that children actually spent more time in close proximity to the stove after it had been fixed than before. In all likelihood 'fixed' stoves still emit dangerous pollutants and spending more time close to it may work against the intention of the intervention. Even though the promotion of stove fixing behaviours has the greatest potential for improving levels of air quality, they may not be feasible because of the cost and the potential risk that children may actually spend more time close to the stove once it is fixed.

Although reducing the duration of burning had a relatively high acceptability rating (26 out of 27 families agreed to try it), it proved to be difficult for most households, not only because of the need for warmth generated by the stove, but also because of the increased burden placed on mothers to perform all of their domestic activities during a shorter period of time. The fact that families need and use the fire for a prolonged period of time makes reducing the duration of solid fuel burning impractical. Even though those families (n=13) that did manage to reduce their burning did so by 13-19%, reducing the duration of burning may have limited feasibility in this context.

It is recommended that the intervention should focus on improving child location and ventilation practices, as they appear to be the most acceptable and feasible in terms of behaviour change. Not only were most families willing to perform the behaviours but they were also able to improve these practices in a measurable way. In terms of moving the child away from the burning fire, the intervention should recommend that someone, preferably an adult or adolescent, watch and care for the child in a location where the smoke from the fire cannot reach. In homes where there is no extra help for mothers children should be kept in the kitchen but as far away from the fire as possible.

The intervention should also focus on encouraging families to open two sources of ventilation during burning, particularly during periods of high smoke emissions such as during ignition or when fuel is added to the fire. The cost of fixing broken windows may serve as a barrier to ventilation practices. Nonetheless, most households have a door leading to the outside as well as a working window so the opening of doors as well as windows can be encouraged.

It is also encouraging to note that 83% of families indicated that they intended to continue with the behaviours after the trials. The most common reason cited for wanting to continue was because of perceived improvements in their children's respiratory health. While these figures may also be exaggerated somewhat by what families thought researchers wanted to hear, it nevertheless highlighted, as in phase one, the importance of mothers' intentions to care for and nurture their children's health as a key motivation for sustained behaviour change. Open-ended interviews showed how other motivations, such as having a cleaner house and saving money through using less fuel, did not feature as strongly as the concern over their children's health. Researchers observed that many mothers viewed smoke as an irritant associated with symptoms such as coughing and eye irritation but were not aware of the association between smoke and serious illnesses such as pneumonia.

It is recommended that the intervention should focus on consolidating mothers' existing knowledge of indoor air pollution but also expand that knowledge to include the dangers of smoke that mothers are not generally aware of such as the dangers of non-visible smoke. The key motivation should be the protection of their families', particularly their children's, respiratory health.

7. REFERENCES

- Albalak, R., Bruce, N., McCracken, J. P., Smith, K. R., & De Gallardo, T. (2001). Indoor respirable particulate matter concentrations from an open fire, improved cookstove and LPG/open fire combination in a rural Guatemalan community. Environ Sci Technol, *35*(13), 1650-5.
- Ballard-Tremmer, G. & Mathee, A. (May, 2000). Review of interventions to reduce the exposure of women and young children to indoor air pollution in developing countries. Paper presented at the USAID and WHO Global Consultation, Health Impacts of indoor air pollution and household energy in developing countries. Washington D.C.
- Barnes, B. R. & Mathee, A. (2002). Reducing childhood exposure to indoor air pollution: the potential role of behaviour change interventions. The Clean Air Journal, *11*(1), 14-18.
- Barnes, B. R. & Mathee, A. (2002). The identification of behavioural intervention opportunities to reduce child exposure to indoor air pollution in rural South Africa. Johannesburg: MRC
- Bruce, N., Perez-Padilla, R., Albalak, R. (2000) Indoor air pollution in developing countries: a major environmental and public health challenge. Bulletin of the World Health Organisation, *78* (9), 1078-1098.
- Cave, B. & Curtis, C. (1999). The effectiveness of promotional techniques in environmental health. WELL study task no: 165. Loughborough University.
- Dickin, K. & Griffiths, M. (1997). Designing by dialogue: a program planners guide to consultative research for improving young child feeding. Report for the Health and Human Resources Analysis Project.
- Ezzati, M. & Kammen, D. M. (2002). The health impacts of exposure to indoor air pollution from solid fuels in developing countries: knowledge, gaps, data needs and policy options. Environmental Health Perspectives, *110*(11), 1058-1068.
- Ezzati, M. & Kammen, D. (2002). Evaluating the health benefits of transitions in household energy technology in Kenya. Energy Policy, *30* (10), 815-26.
- Ezzati, M., Saleh, H., & Kamen, D. M. (2000). The contributions of emissions and spatial micro-environments to exposure to indoor air pollution from biomass combustion in Kenya. Environmental Health Perspectives, *108*(9), 833-9.

- Favin, M., Yacoob, M., & Bendahmane, D. (1999). Behavior first: a minimum package of environmental health behaviors to improve child health. Applied study no. 10. Washington D. C.: Environmental Health Project.
- Gitonga, S. (2001). Smoke project activities. Unpublished project report. Kenya: ITDG
- Miles, M. B., & Huberman, A. M. (1994). Qualitative data analysis (2 ed.). Thousand Oaks: Sage.
- Naeher, L. P., Leaderer, B. P., & Smith, K. R. (2000). Particulate matter and carbon monoxide in highland Guatemala: indoor and outdoor levels from traditional and improved wood stoves and gas stoves. Indoor Air, 10(3), 200-5.
- Patton, M. Q. (1990). Qualitative evaluation and research methods. Thousand Oaks: Sage
- Raphael, D. (2000). The question of evidence in health promotion. Health Promotion International, 15 (4), p. 355-367.
- Smith, K. R. (1999). Indoor air pollution: the global problem. Pollution management in focus, discussion note number 4. Geneva: The World Bank.
- Smith, K. R. (1987). Biofuels, air pollution and health. New York: Plenum Press.
- Smith, K. R., Samet, J. M., Romieu, I., & Bruce, N. (2000). Indoor air pollution in developing countries and acute lower respiratory infection in children. Thorax, 55, 518-532.
- The World Resources Institute. (1998). A guide to the global environment. Oxford: Oxford University Press.
- Von Schirnding, Y., Bruce, N., Smith, K. R., Ballard-Tremeer, G., Ezzati, M., Lvovsky, K. (2002). Addressing the impact of household energy and indoor air pollution on the health of the poor. Geneva: World Health Organisation.
- Wafula, E. M., Kinyanjui, M. M., Nyabola, L., & Tenambergen, E. D. (2000). Effect of improved stoves on prevalence of acute respiratory infection and conjunctivitis among children and women in a rural community in Kenya. East African Medical Journal, 77(1), 37-41.

ⁱ Strong evidence shows that exposure to indoor air pollution is also associated with an increased risk of chronic obstructive lung disease (COLD) in adults and moderate to weak evidence exists for Tuberculosis, adverse birth outcomes, eye problems and cardiovascular disease.